



Colorectal Cancer Fact Sheet

for Patients and Caregivers



Colorectal cancer starts in either the colon or rectum. In the US, colorectal cancer is the third most common cancer. It is also the second-leading cause of cancer death in both men and women. Most colorectal cancers start with a polyp. A polyp is a small growth on the lining of the colon or rectum. Not all polyps develop into cancer.

Risk Factors

More than half of colorectal cancers can be linked to risk factors that can be changed. These include excess body weight, not being physically active, smoking, and drinking alcohol. Risk can also be increased by eating a lot of red or processed meats, or by not eating enough fruits, vegetables, and whole grains.

There are other risk factors that can't be changed. These include: your age; having a personal or family history of colorectal polyps; having inflammatory bowel disease, certain hereditary syndromes (such as Lynch syndrome), or type 2 diabetes; and your race and ethnicity. Black people, American Indian and Alaska Native people, and people of Ashkenazi Jewish descent are at higher risk for colorectal cancer.

If you have certain risk factors, your healthcare provider may have you see a genetic counselor. They can help you better understand your risks and make informed decisions about genetic testing and your care.

Prevention

Not all colorectal cancers can be prevented. But you can help lower your risk.

- Get regular screening to help find problems that might turn into cancer, such as polyps that can be removed.
- Get to and stay at a healthy weight.
- Be as active as you can be.

- Eat a healthy diet with plenty of fruits, vegetables, and whole grains. Limit or avoid red and processed meats and highly processed foods.
- Avoid alcohol, or limit yourself to 1 drink per day for women or 2 drinks per day for men.
- Avoid tobacco, and quit if you smoke.

Screening and Early Detection

Screening is a process used to look for cancer in people who have no symptoms. Talk to a healthcare provider about your risks for colorectal cancer and when to start screening. Also, talk to your insurance company about whether they pay the cost of screening.

The American Cancer Society recommends the following for people at average risk for colorectal cancer:

- Start regular screening at **age 45**.
- People who are in good health should continue regular colorectal cancer screening through the **age of 75**.
- People **ages 76 through 85** should talk to their healthcare provider about whether to be screened based on their preferences, life expectancy, health status, and past screening history.
- People **over age 85** should no longer get colorectal cancer screening.

People at **high risk** of colorectal cancer based on family or personal history or other factors may need to start screening before age 45, be screened more often, or get specific tests.

Screening for colorectal cancer can be done with different types of tests.

Stool-based tests (preferred)

These tests look for signs of cancer in a person's stool. Options include:

- Fecal immunochemical test (FIT)* every year, or
- Guaiac-based fecal occult blood test (gFOBT)* every year, or
- Multi-targeted stool DNA (mt-sDNA) test* every 3 years, or
- Multi-targeted stool RNA (mt-sRNA) test* every 3 years

Visual exams of the colon and rectum (preferred)

These tests look inside the colon and rectum for any abnormal areas. Options include:

- Colonoscopy every 10 years, or
- CT colonography (virtual colonoscopy)* every 5 years, or
- Sigmoidoscopy* every 5 years

Blood-based tests (not preferred)

Blood-based tests* check for signs of colorectal cancer in a sample of a person's blood. While these tests are an option, they're not preferred at this time. They should only be used if you're not willing to get a stool-based test or visual exam.

**If a person chooses to be screened with a test other than a colonoscopy, any abnormal test result should be followed up with a colonoscopy.*

There are some differences between these tests to consider, but **the most important thing is to get screened, no matter which test you choose.**

Signs and Symptoms

Many colorectal cancers do not cause symptoms right away. But, some signs and symptoms of colorectal cancer might include:

- Bleeding from your rectum
- Blood in your stool
- Change in your bowel habits or stool shape
- Stomach cramping or pain
- Feeling the need to have a bowel movement that doesn't go away after you have one



- Not feeling hungry
- Losing weight without trying
- Weakness

Treatment

Treatment for colorectal cancer depends on several things. These include the type and extent of the cancer, the person's age, other health problems, and personal choices. Also, special testing might be done on the tumor to figure out the best treatment. If you have colorectal cancer, talk to your doctor about the best treatment for you.

Living With Colorectal Cancer

Having colorectal cancer affects a person's quality of life. Physical, social, mental health, spiritual, and money issues can come up at any time.

People with colorectal cancer may also be helped by palliative care at any time. Palliative care focuses on helping manage symptoms, address issues, and improve quality of life.

Good communication between a person with cancer and their healthcare team is important. This should include:

- Asking and answering questions
- Working together to set care goals
- Shared decision-making
- Managing side effects and other issues
- Scheduling follow-up tests and care



For more information and support, visit the American Cancer Society website at cancer.org/colorectalcancer or call us at **1-800-227-2345**. We're here when you need us.



cancer.org | 1.800.227.2345

©2026, American Cancer Society, Inc.
No.080764 Rev. 5/26
Models used for illustrative purposes only.

